

About the Celestial Objects

Listed on this page are several of the brighter, more interesting celestial objects visible in the evening sky this month (refer to the monthly sky map). The objects are grouped into three categories. Those that can be easily seen with the naked eye (that is, without optical aid), those easily seen with binoculars, and those requiring a telescope to be appreciated. **Note, all of the objects (except single stars) will appear more impressive when viewed through a telescope or very large binoculars.** They are grouped in this way to highlight objects that can be seen using the optical equipment that may be available to the star gazer.

Tips for Observing the Night Sky

When observing the night sky, and in particular deep-sky objects such as star clusters, nebulae, and galaxies, it's always best to observe from a dark location. Avoid direct light from street lights and other sources. If possible observe from a dark location away from the light pollution that surrounds many of today's large cities.

You will see more stars after your eyes adapt to the darkness—usually about 10 to 20 minutes after you go outside. Also, if you need to use a torch to view the sky map, cover the light bulb with red cellophane. This will preserve your dark vision.

Finally, even though the Moon is one of the most stunning objects to view through a telescope, its light is so bright that it brightens the sky and makes many of the fainter objects very difficult to see. So try to observe the evening sky on moonless nights around either New Moon or Last Quarter.

Astronomical Glossary

Conjunction – An alignment of two celestial bodies such that they present the least angular separation as viewed from Earth.

Constellation – A defined area of the sky containing a star pattern.

Diffuse Nebula – A cloud of gas illuminated by nearby stars.

Double Star – Two stars that appear close to each other in the sky; either linked by gravity so that they orbit each other (binary star) or lying at different distances from Earth (optical double). Apparent separation of stars is given in seconds of arc (").

Ecliptic – The path of the Sun's center on the celestial sphere as seen from Earth.

Elongation – The angular separation of two celestial bodies. For Mercury and Venus the greatest elongation occurs when they are at their most angular distance from the Sun as viewed from Earth.

Galaxy – A mass of up to several billion stars held together by gravity.

Globular Star Cluster – A ball-shaped group of several thousand old stars.

Light Year (ly) – The distance a beam of light travels at 300,000 km/sec in one year.

Magnitude – The brightness of a celestial object as it appears in the sky.

Open Star Cluster – A group of tens or hundreds of relatively young stars.

Opposition – When a celestial body is opposite the Sun in the sky.

Planetary Nebula – The remnants of a shell of gas blown off by a star.

Universal Time (UT, UTC, or GMT) – A time system used by astronomers. USA Eastern Standard Time (for example, New York) is 5 hours behind 0 hours UT.

Variable Star - A star that changes brightness over a period of time.

NORTHERN HEMISPHERE MAY 2007

OBJE(

LESTIAL (

Easily Seen with the Naked Eye

Capella	Aur	• The 6th brightest star. Appears yellowish in color. Spectroscopic binary. Dist=42 ly.
Arcturus	Boo	 Orange, giant K star. Name means "bear watcher". Dist=37 ly.
Procyon	CMi	• Greek name meaning "before the dog" - rises before Sirius (northern latitudes). Dist=11.4 ly.
δ Cephei	Сер	Cepheid prototype. Mag varies between 3.5 & 4.4 over 5.366 days. Mag 6 companion.
Deneb	Cyg	 Brightest star in Cygnus. One of the greatest known supergiants. Dist=3,000 ly.
Castor	Gem	 Multiple star system with 6 components. 3 stars visible in telescope. Dist=52 ly.
Pollux	Gem	 With Castor, the twin sons of Leda in classical mythology. Dist=34 ly.
α Herculis	Her	Semi-regular variable. Magnitude varies between 3.1 & 3.9 over 90 days. Mag 5.4 companion.
Regulus	Leo	 Brightest star in Leo. A blue-white star with at least 1 companion. Dist=77 ly.
Vega	Lyr	 The 5th brightest star in the sky. A blue-white star. Dist=25.3 ly.
Antares	Sco	 Red, supergiant star. Name means "rival of Mars". Dist=135.9 ly.
Polaris	UMi	 The North Pole Star. A telescope reveals an unrelated mag 8 companion star.
Spica	Vir	• Latin name means "ear of wheat" and shown held in Virgo's left hand. Dist=260 ly.

Easily Seen with Binoculars

M44	Cnc	0	Praesepe or Beehive Cluster. Visible to the naked eye. Dist=577 ly.
M3	CVn	⊕	Easy to find in binoculars. Might be glimpsed with the naked eye.
μ Cephei	Сер	•	Herschel's Garnet Star. One of the reddest stars. Mag 3.4 to 5.1 over 730 days.
Mel 111	Com	ं	Coma Berenices. 80 mag 5-6 stars in 5 deq. Dist=288 ly. Age=400 million years.
χ Cyqni		•	Long period pulsating red giant. Magnitude varies between 3.3 & 14.2 over 407 days.
,, ,,	Cyg		31 1 3 3 3
M39	Cyg	0	May be visible to the naked eye under good conditions. Dist=900 ly.
v Draconis	Dra	•	Wide pair of white stars. One of the finest binocular pairs in the sky. Dist=100 ly.
M13	Her	Ф	Best globular in northern skies. Discovered by Halley in 1714. Dist=23,000 ly.
M92	Her	0	Fainter and smaller than M13. Use a telescope to resolve its stars.
R Hydrae	Hya	•	Long period variable. Mag varies between 3.0 & 11.0 over 390 days. Brilliant red.
ε Lyrae	Lyr	•	Famous Double Double. Binoculars show a double star. High power reveals each a double.
R Lyrae	Lyr	•	Semi-regular variable. Magnitude varies between 3.9 & 5.0 over 46.0 days.
M12	0ph	Ф	Close to the brighter M10. Dist=18,000 ly.
M10	0ph	Ф	3 degrees from the fainter M12. Both may be glimpsed in binoculars. Dist=14,000 ly.
IC 4665	0ph	0	Large, scattered open cluster. Visible with binoculars.
6633	0ph	0	Scattered open cluster. Visible with binoculars.
M4	Sco	0	A close globular. May just be visible without optical aid. Dist=7,000 ly.
M5	Ser	0	Fine globular star cluster. Telescope will reveal individual stars. Dist=25,000 ly.
Mizar & Alcor	UMa	•	Good eyesight or binoculars reveals 2 stars. Not a binary. Mizar has a mag 4 companion.
Cr 399	Vul	0	Coathanger asterism or "Brocchi's Cluster". Not a true star cluster. Dist=218 to 1,140 ly.

Telescopic Objects

JICSCOP	,,,,	,,,	,0013
oötis	Boo	•	Red giant star (mag 2.5) with a blue-green mag 4.9 companion. Sep=2.8". Difficult to split.
7	Cnc	\Diamond	Contains 500+ stars mag 10 & fainter. One of the oldest clusters. Dist=2,350 ly.
4	CVn	0	Compact nearly face-on spiral galaxy. Dist=15 million ly.
assiopeiae	Cas	•	Yellow star mag 3.4 & orange star mag 7.5. Dist=19 ly. Orbit=480 years. Sep=12".
28	Cen	0	Bisected by a wide obscuring lane. Strong radio source. Dist=14 million ly.
1	CVn	0	Whirlpool Galaxy. First recognised to have spiral structure. Dist=25 million ly.
4	Com	0	Black-Eye Galaxy. Discovered by J.E. Bode in 1775 - "a small, nebulous star".
ireo	Cyg	•	Beautiful double star. Contrasting colours of orange and blue-green. Sep=34.4".
Cygni	Cyg	•	Attractive double star. Mags 5.2 & 6.1 orange dwarfs. Dist=11.4 ly. Sep=28.4".
12	Hya		Ghost of Jupiter. Bright blue disk. Mag 11 central star. Dist=2,600 ly.
3	Hya	0	Classic face-on spiral. Discovered in 1752 by Lacaille. In attractive star field.
eonis	Leo	•	Superb pair of golden-yellow giant stars. Mags 2.2 & 3.5. Orbit=600 years. Sep=4.4".
yrae	Lyr	•	Eclipsing binary. Mag varies between 3.3 & 4.3 over 12.940 days. Fainter mag 7.2 blue star.
7	Lyr		Ring Nebula. Magnificent object. Smoke-ring shape. Dist=4,100 ly.
6	Ser		Eagle Nebula. Requires a telescope of large aperture. Dist=8,150 ly.
1	UMa	0	Beautiful spiral galaxy visible with binoculars. Easy to see in a telescope.
2	UMa	0	Close to M81 but much fainter and smaller.
04	Vir	0	Sombrero Galaxy. Almost edge-on spiral galaxy. Protruding central core.
irginis	Vir	•	Superb pair of mag 3.5 yellow-white stars. Orbit=169 years. At their closest in 2005.
	oötis 7 4 assiopeiae 8 1 4 ireo Cygni 2 3 eonis yrae 7 5 1 1 2 0 4	oötis Boo Character Company Co	7 Cnc

Dumbbell Nebula. Large, twin-lobed shape. Most spectacular planetary. Dist=975 ly.

M27